

The bounce-splash of a viscoelastic drop

Federico Hernández-Sánchez, René Ledesma & Roberto Zenit

Instituto de Investigaciones en Materiales
Universidad Nacional Autónoma de México
Cd. Universitaria, México D.F., MÉXICO

October 10, 2008

Abstract

This is an entry for the Gallery of Fluid Motion of the 61st Annual Meeting of the APS-DFD (fluid dynamics videos). This video shows the collision and rebound of viscoelastic drops against a solid wall. Using a high speed camera, the process of approach, contact and rebound of drops of a viscoelastic liquid is observed. We found that these drops first splash, similar to what is observed in Newtonian colliding drops; after a few instants, the liquid recoils, recovering its original drop shape and bounce off the wall.

1 Introduction

We are interested in studying non-Newtonian phenomena in general; in particular, we have studied the motion of spheres and bubbles in complex fluids[1, 2]. We also have previously studied the collision of solid spheres immersed in viscous liquids[3] and more recently the collision of bubbles[4]. This visualization project shows the continuation of our previous work in these two subjects.

2 Experimental Conditions

We release viscoelastic drops of approximately 2.2 cm in diameter from different heights in air to produce collisions at various speeds against a solid surface. Different aqueous mixtures of gelatin were used as the test liquid. By varying the viscoelastic properties of the drops and the collision

speed we can change the value of the relevant dimensionless number. The Weissenberg number is defined as

$$We = \frac{\lambda U}{D} \quad (1)$$

where λ is the relaxation time, U is the impact velocity and D is the drop diameter. Hence, a large value of We would show a liquid-like behavior; on the other hand, a small We collision would resemble a solid rebound. In this video contribution the collision process of drops for a range of values of Weissenberg numbers is shown.

The most interesting case shown is that for the largest We . At contact, the drops splash: the whole liquid volume spreads into a thin sheet. After a few instants, as a result of the elasticity of the media, the liquid sheet recoils. The drop is re-formed, but highly distorted. Nevertheless, the drop bounces off the wall and exhibits large shape oscillations.

To our knowledge this process has not been studied in the past.

3 Videos

Our video contributions can be found at:

- Video 1, mpeg2, full resolution
- Video 2, mpeg1, low resolution

References

- [1] Soto, E., Goujon, C., Zenit, R. and Manero, O. “A study of velocity discontinuity for single air bubbles rising in an associative polymer,” *Phys. Fluids*, **18**, 121510,(2006)
- [2] A.J. Mendoza-Fuentes, R. Montiel, R. Zenit, O. Manero, “On the flow of associative polymers past a sphere: Evaluation of negative wake criteria,” submitted to *Phys. Fluids*, (2008).
- [3] G. G. Joseph, R. Zenit, M. L. Hunt, and A. M. Rosenwinkel, “Particle wall collision in a viscous fluid,” *J. Fluid Mech.* **433**, 329 (2001).
- [4] R. Zenit and D. Legendre, “The coefficient of restitution for air bubbles colliding against solid walls in viscous liquids,” submitted to *Phys. Fluids*, (2008).